

Please check the examination details below before entering your candidate information

Candidate surname					Other names			
Pearson Edexcel		Centre Number			Candidate Number			
		<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
Tuesday 2 June 2020								
Afternoon (Time: 1 hour 45 minutes)					Paper Reference 9CH0/01			
Chemistry								
Advanced								
Paper 1: Advanced Inorganic and Physical Chemistry								
Candidates must have: Scientific calculator							Total Marks	
Data Booklet								
Ruler								

Instructions

- Use **black** ink or **black** ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided
– *there may be more space than you need.*

Information

- The total mark for this paper is 90.
- The marks for **each** question are shown in brackets
– *use this as a guide as to how much time to spend on each question.*
- For the question marked with an **asterisk** (*), marks will be awarded for your ability to structure your answer logically showing the points that you make are related or follow on from each other where appropriate.
- A Periodic Table is printed on the back cover of this paper.

Advice

- Read each question carefully before you start to answer it.
- Show all your working in calculations and include units where appropriate.
- Check your answers if you have time at the end.

Turn over ►

P62668A

©2020 Pearson Education Ltd.

1/1/1/1/1



Pearson

Answer ALL questions.

Some questions must be answered with a cross in a box ☒. If you change your mind about an answer, put a line through the box ☒ and then mark your new answer with a cross ☒.

1 (a) Which equation shows the third ionisation energy of aluminium?

(1)

- A** $\text{Al(g)} \rightarrow \text{Al}^{3+}(\text{g}) + 3\text{e}^{-}$
- B** $\text{Al}^{2+}(\text{g}) \rightarrow \text{Al}^{3+}(\text{g}) + \text{e}^{-}$
- C** $\text{Al}^{3+}(\text{g}) + 3\text{e}^{-} \rightarrow \text{Al(g)}$
- D** $\text{Al}^{3+}(\text{g}) + \text{e}^{-} \rightarrow \text{Al}^{2+}(\text{g})$

(b) Which element in this table is in Group 2?

Element	Ionisation energy / kJ mol^{-1}			
	First	Second	Third	Fourth
W	1086	2353	4621	6223
X	653	1592	2987	4740
Y	590	1145	4912	6474
Z	496	4563	6913	9544


(1)

- A** W
- B** X
- C** Y
- D** Z



- (c) The graph shows the first ionisation energies (IE) of eight successive elements from the first 20 elements in the Periodic Table.

Which letter represents the first ionisation energy of oxygen?



(1)

- A** (a)
- B** (b)
- C** (c)
- D** (h)

- (d) Give the formula of a stable **ion** that is isoelectronic with the magnesium ion, Mg^{2+} .

(1)



(e) A student stated that 'the elements scandium and zinc are d-block elements but are not transition metals'.

Discuss this statement, using appropriate electronic configurations to support your answer.

(4)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(Total for Question 1 = 8 marks)



2 This question is about acids and bases.

(a) What is the order of **decreasing** pH for $0.100 \text{ mol dm}^{-3}$ solutions of these three acids? (1)

- A $\text{CH}_3\text{COOH} > \text{CH}_2\text{ClCOOH} > \text{HCl}$
- B $\text{HCl} > \text{CH}_3\text{COOH} > \text{CH}_2\text{ClCOOH}$
- C $\text{CH}_2\text{ClCOOH} > \text{CH}_3\text{COOH} > \text{HCl}$
- D $\text{HCl} > \text{CH}_2\text{ClCOOH} > \text{CH}_3\text{COOH}$

(b) A solution of methanoic acid, HCOOH , has a concentration of $0.240 \text{ mol dm}^{-3}$ and a pH of 2.20.

Calculate the value of $\text{p}K_{\text{a}}$ for methanoic acid.

(3)

(c) Which of these mixtures would form a buffer solution with a pH **below** 7? (1)

- A NaOH(aq) and excess HCl(aq)
- B NaOH(aq) and excess $\text{CH}_3\text{COOH(aq)}$
- C excess NaOH(aq) and HCl(aq)
- D excess NaOH(aq) and $\text{CH}_3\text{COOH(aq)}$



(d) Bromothymol blue, methyl orange and phenolphthalein are indicators used in titrations.

Which, if any, of these indicators could be used for a titration of ammonia, $\text{NH}_3(\text{aq})$, with ethanoic acid, $\text{CH}_3\text{COOH}(\text{aq})$?

(1)

- A** bromothymol blue
- B** methyl orange
- C** phenolphthalein
- D** none of these three indicators

(Total for Question 2 = 6 marks)



3 This question is about transition metals and transition metal complexes.

- (a) Describe the bonding in the element chromium and use your answer to justify why it has such a high melting temperature.

You may find it helpful to draw a labelled diagram.

(4)

.....

.....

.....

.....

.....

.....

.....

- (b) When chromium(III) sulfate dissolves in water, a green solution containing the $[\text{Cr}(\text{H}_2\text{O})_6]^{3+}$ ion forms.

(i) Give the shape of this complex ion.

(1)

.....

(ii) Explain why the chromium complex ion is coloured.

(3)

.....

.....

.....

.....

.....

.....

.....


.....

.....


.....



(c) The ligand ethylenediaminetetraacetate, EDTA^{4-} , has the structure shown.



When a solution of EDTA^{4-} is added to a solution of $[\text{Cr}(\text{H}_2\text{O})_6]^{3+}$ ions, a new complex ion is formed.



The equilibrium constant for this equilibrium is $2.51 \times 10^{23} \text{ dm}^3 \text{ mol}^{-1}$.

By considering the equilibrium for this reaction and changes in entropy, comment on the value of the equilibrium constant. No calculations are required.


(3)



- (d) Aqueous vanadium(II) chloride, $\text{VCl}_2(\text{aq})$, can be oxidised by bubbling gaseous chlorine, $\text{Cl}_2(\text{g})$, through the solution in the absence of air.

40.0 cm^3 of $0.100 \text{ mol dm}^{-3}$ VCl_2 solution was oxidised by 144 cm^3 of chlorine gas, at room temperature and pressure (r.t.p.).

The chlorine was reduced to chloride ions, according to the half-equation



[Molar volume of a gas at r.t.p. = $24.0 \text{ dm}^3 \text{ mol}^{-1}$]

- (i) Use these data to calculate the final oxidation state of vanadium.
You **must** show your working.

(5)

- (ii) State the initial and final colours you would see as the chlorine bubbles through the aqueous vanadium(II) chloride, $\text{VCl}_2(\text{aq})$.

(2)

(Total for Question 3 = 18 marks)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

BLANK PAGE




4 This question is about dissolving different compounds.

(a) Which of these compounds is the most soluble in water?

(1)

- A** barium sulfate
- B** calcium sulfate
- C** magnesium sulfate
- D** strontium sulfate

(b) What is the value, in kJ mol^{-1} , for the standard enthalpy change of solution of magnesium chloride?



$$\text{Lattice energy MgCl}_2(\text{s}) = -2526 \text{ kJ mol}^{-1}$$

$$\text{Hydration enthalpy of Cl}^{-}(\text{g}) = -381 \text{ kJ mol}^{-1}$$

$$\text{Hydration enthalpy of Mg}^{2+}(\text{g}) = -1921 \text{ kJ mol}^{-1}$$

(1)

- A** +157
- B** -157
- C** +224
- D** -224



* (c) The solubility of two compounds in different solvents was investigated.
A summary of the findings is shown.

Compound	Soluble in water	Soluble in hexane
2-methylpentane	X	✓
potassium bromide	✓	X

Explain the findings of the investigation by considering the interactions between the compounds and each of the solvents.

(6)

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Large area with horizontal dotted lines for writing.

(Total for Question 4 = 8 marks)



5 This question is about the chemistry of hydrated magnesium nitrate, $\text{Mg}(\text{NO}_3)_2 \cdot x\text{H}_2\text{O}$.

(a) Group 2 nitrates decompose when heated.

(i) State **two** observations you would see when hydrated magnesium nitrate is heated. (2)

.....

.....

.....

(ii) Explain the trend in thermal stability of Group 2 nitrates. (3)

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) In an experiment, a sample of hydrated magnesium nitrate, $\text{Mg}(\text{NO}_3)_2 \cdot x\text{H}_2\text{O}$, with a mass of 0.765 g, was dissolved in water and reacted with an excess of sodium hydroxide solution, $\text{NaOH}(\text{aq})$. The precipitate of magnesium hydroxide, $\text{Mg}(\text{OH})_2$, produced was removed and dried. The mass of the dried sample was 0.174 g.

(i) Draw dot-and-cross diagrams for the ions in magnesium hydroxide. Show the outer electrons only. (2)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



- (ii) Use the experimental data to calculate the value for x in the formula $\text{Mg}(\text{NO}_3)_2 \cdot x\text{H}_2\text{O}$.
You **must** show all your working.

(5)

DO NOT WRITE IN THIS AREA


DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(Total for Question 5 = 12 marks)




6 Prop-2-en-1-ol is an unsaturated alcohol with the structure shown.



- (a) A student planned to use bond enthalpy data to calculate a value for the enthalpy change of combustion of prop-2-en-1-ol.
- (i) When researching the bond enthalpy data, the student claimed that it was not necessary to find the value for the C=C bond as they could use the value for a C–C bond and multiply it by two. Explain why the student is **incorrect**.

(2)

- (ii) Calculate a value for the enthalpy of combustion of prop-2-en-1-ol using the data shown.



Bond	C–C	C=C	C–O	C=O	O–H	C–H	O=O
Bond enthalpy / kJ mol^{-1}	347	612	358	805	464	413	498

(3)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



(iii) Explain, in terms of entropy, why the combustion of prop-2-en-1-ol is always feasible in the gaseous state.

(2)


.....

.....

.....

.....

(b) Chemists are researching a process to make ethanol and ethene directly from carbon dioxide and water.



	$\text{CO}_2(\text{g})$	$\text{H}_2\text{O}(\text{l})$	$\text{CH}_3\text{CH}_2\text{OH}(\text{l})$	$\text{C}_2\text{H}_4(\text{g})$	$\text{O}_2(\text{g})$
$S^\ominus / \text{J K}^{-1} \text{ mol}^{-1}$	213.6	69.9	160.7	219.5	205.0


Calculate $\Delta S^\ominus_{\text{total}}$ for the reaction and hence determine whether the reaction is feasible under standard conditions.

(5)

(Total for Question 6 = 12 marks)



- 7 A mixture of ethanoic acid, ethanol and a catalyst was left for several days to reach equilibrium.



The equilibrium constant, K_c , **under these conditions**, was 0.28.

- (a) (i) Write the expression for the equilibrium constant, K_c .

(1)

- (ii) The initial amounts of ethanol and ethanoic acid used were 1.2 mol of each reactant.

Use this information, your expression for the equilibrium constant, K_c , and the value for K_c , to find the amounts of each product at equilibrium, in moles.

(3)

Amount of $\text{CH}_3\text{COOCH}_2\text{CH}_3$ =

Amount of H_2O =


DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



- (b) Another ester, methyl methanoate, can be formed by the reaction between methanol and carbon monoxide in the gaseous phase.



- (i) The two O–C–H bond angles, x and y, in the ester are approximately (1)

- A 180° and 90°
 B 120° and 90°
 C 120° and 109.5°
 D 109.5° and 109.5°

- (ii) The reaction often forms an equilibrium mixture.

Which could be the units for the equilibrium constant, K_p ?

(1)

- A mol dm⁻³
 B dm³ mol⁻¹
 C atm
 D atm⁻¹

- (iii) Describe what effect, if any, increasing the pressure would have on the equilibrium constant, K_p . Justify your answer.

(2)

.....

.....

.....

.....

.....

.....

.....

.....

.....

(Total for Question 7 = 8 marks)




- 8 Tablets containing potassium manganate(VII), KMnO_4 , are dissolved in water forming an antiseptic solution to treat skin conditions. The manufacturers claim that each tablet contains 400 mg of KMnO_4 .

To check the claim, the titration procedure outlined was carried out.


- Five tablets were dissolved in distilled water to make 100.0 cm^3 of solution.
- Some of the KMnO_4 solution was used to fill a burette.
- 25.0 cm^3 of sodium ethanedioate solution, $\text{Na}_2\text{C}_2\text{O}_4(\text{aq})$, of concentration $0.200 \text{ mol dm}^{-3}$, was added to a conical flask and warmed.
- Sulfuric acid, of concentration 2 mol dm^{-3} , was also added to the conical flask.
- The KMnO_4 solution was added to the flask from the burette, until the end-point.

The equation for the reaction between MnO_4^- ions from the KMnO_4 and $\text{C}_2\text{O}_4^{2-}$ ions from the sodium ethanedioate solution is shown.



- (a) Give the colour **change** at the end-point of the titration. (1)

- (b) (i) Complete the dot-and-cross diagram for the ethanedioate ion.
Show the outer electrons only. (2)



- (ii) Determine the oxidation number of carbon in the ethanedioate ion, $\text{C}_2\text{O}_4^{2-}$. (1)





(c) Give the reason why sulfuric acid was also added to the conical flask.

(1)

(d) This redox reaction could be used in an electrochemical cell.

The cell half-equations are



Write a cell diagram for this cell using the conventional representation.

(2)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA




(e) The results of the titration are shown.

Run	Trial	1	2	3
Final volume / cm ³	17.50	34.10	17.20	34.10
Initial volume / cm ³	0.00	17.30	0.00	17.20
Titre / cm ³	17.50		17.20	
Concordant titres (✓)				
Mean titre / cm ³				

(i) Complete the table.

(2)

(ii) The equation for the reaction between MnO_4^- ions from the KMnO_4 and $\text{C}_2\text{O}_4^{2-}$ ions from the sodium ethanedioate solution is shown.



Use this equation and your mean titre from (e)(i) to calculate the mass, in mg, of KMnO_4 in **one** tablet.

Give your answer to an appropriate number of significant figures.

(5)



- (iii) A textbook suggested the conical flask should be heated during the titration, as the reaction between the MnO_4^- ions and the $\text{C}_2\text{O}_4^{2-}$ ions is slow.

Use these electrode potentials and your knowledge of homogeneous catalysis to deduce why the heating is very important at the start of the titration, but less important as the titration proceeds. Justify your answer. You may include equations in your justification.

Electrode system	E^\ominus / V
$2\text{CO}_2(\text{g}) + 2\text{e}^- \rightleftharpoons \text{C}_2\text{O}_4^{2-}(\text{aq})$	+0.64
$\text{Mn}^{3+}(\text{aq}) + \text{e}^- \rightleftharpoons \text{Mn}^{2+}(\text{aq})$	+1.49
$\text{MnO}_4^-(\text{aq}) + 8\text{H}^+(\text{aq}) + 5\text{e}^- \rightleftharpoons \text{Mn}^{2+}(\text{aq}) + 4\text{H}_2\text{O}(\text{l})$	+1.51

(4)

(Total for Question 8 = 18 marks)

TOTAL FOR PAPER = 90 MARKS

The Periodic Table of Elements

		1	2											3	4	5	6	7	0 (8)	
		(2)												(18)						
(1)	6.9 Li lithium 3	9.0 Be beryllium 4											10.8 B boron 5	12.0 C carbon 6	14.0 N nitrogen 7	16.0 O oxygen 8	19.0 F fluorine 9	20.2 Ne neon 10		
	23.0 Na sodium 11	24.3 Mg magnesium 12											27.0 Al aluminium 13	28.1 Si silicon 14	31.0 P phosphorus 15	32.1 S sulfur 16	35.5 Cl chlorine 17	39.9 Ar argon 18		
	39.1 K potassium 19	40.1 Ca calcium 20	45.0 Sc scandium 21	47.9 Ti titanium 22	50.9 V vanadium 23	52.0 Cr chromium 24	54.9 Mn manganese 25	55.8 Fe iron 26	58.9 Co cobalt 27	58.7 Ni nickel 28	63.5 Cu copper 29	65.4 Zn zinc 30	69.7 Ga gallium 31	72.6 Ge germanium 32	74.9 As arsenic 33	79.0 Se selenium 34	79.9 Br bromine 35	83.8 Kr krypton 36		
	85.5 Rb rubidium 37	87.6 Sr strontium 38	88.9 Y yttrium 39	91.2 Zr zirconium 40	92.9 Nb niobium 41	95.9 Mo molybdenum 42	[98] Tc technetium 43	101.1 Ru ruthenium 44	102.9 Rh rhodium 45	106.4 Pd palladium 46	107.9 Ag silver 47	112.4 Cd cadmium 48	114.8 In indium 49	118.7 Sn tin 50	121.8 Sb antimony 51	127.6 Te tellurium 52	126.9 I iodine 53	131.3 Xe xenon 54		
	132.9 Cs caesium 55	137.3 Ba barium 56	138.9 La* lanthanum 57	178.5 Hf hafnium 72	180.9 Ta tantalum 73	183.8 W tungsten 74	186.2 Re rhenium 75	190.2 Os osmium 76	192.2 Ir iridium 77	195.1 Pt platinum 78	197.0 Au gold 79	200.6 Hg mercury 80	204.4 Tl thallium 81	207.2 Pb lead 82	209.0 Bi bismuth 83	[209] Po polonium 84	[210] At astatine 85	[222] Rn radon 86		
	[223] Fr francium 87	[226] Ra radium 88	[227] Ac* actinium 89	[261] Rf rutherfordium 104	[262] Db dubnium 105	[266] Sg seaborgium 106	[264] Bh bohrium 107	[277] Hs hassium 108	[268] Mt meitnerium 109	[271] Ds darmstadtium 110	[272] Rg roentgenium 111	Elements with atomic numbers 112-116 have been reported but not fully authenticated								
	140 Ce cerium 58	141 Pr praseodymium 59	144 Nd neodymium 60	147 Pm promethium 61	150 Sm samarium 62	152 Eu europium 63	157 Gd gadolinium 64	159 Tb terbium 65	163 Dy dysprosium 66	165 Ho holmium 67	167 Er erbium 68	169 Tm thulium 69	173 Yb ytterbium 70	175 Lu lutetium 71						
	232 Th thorium 90	[231] Pa protactinium 91	238 U uranium 92	[237] Np neptunium 93	[242] Pu plutonium 94	[243] Am americium 95	[247] Cm curium 96	[245] Bk berkelium 97	[251] Cf californium 98	[254] Es einsteinium 99	[253] Fm fermium 100	[256] Md mendelevium 101	[254] No nobelium 102	[257] Lr lawrencium 103						

* Lanthanide series
* Actinide series

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



P 6 2 6 6 8 A 0 2 4 2 4