Please check the examination details belo	ow before entering your candidate information
Candidate surname	Other names
Pearson Edexcel Level 3 GCE	tre Number Candidate Number
Monday 18 May	y 2020
Morning (Time: 1 hour 30 minutes)	Paper Reference 8CH0/01
Chemistry Advanced Subsidiary Paper 1: Core Inorganic and	d Physical Chemistry
Candidates must have: Scientific cal Data Bookle Ruler	

Instructions

- Use **black** ink or **black** ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided
 - there may be more space than you need.

Information

- The total mark for this paper is 80.
- The marks for **each** question are shown in brackets
 - use this as a guide as to how much time to spend on each question.
- For the question marked with an asterisk (*), marks will be awarded for your ability to structure your answer logically, showing the points that you make are related or follow on from each other where appropriate.
- A Periodic Table is printed on the back cover of this paper.

Advice

- Read each question carefully before you start to answer it.
- Show all your working in calculations and include units where appropriate.
- Check your answers if you have time at the end.

Turn over ▶







Answer ALL questions.

Some questions must be answered with a cross in a box \boxtimes . If you change your mind about an answer, put a line through the box 🔀 and then mark your new answer with a cross \bowtie .

- This question is about the electronic structure of some Group 5 elements.
 - (a) Which is the electronic configuration of the arsenide ion, As³-?

(1)

- \square **A** $1s^22s^22p^63s^23p^63d^{10}4s^2$
- \square **B** $1s^22s^22p^63s^23p^63d^{10}4s^24p^3$
- \square **C** $1s^22s^22p^63s^23p^63d^{10}4s^24p^6$
- \square **D** 1s²2s²2p⁶3s²3p⁶3d¹⁰4s²4p³4d³
- (b) The electronic configuration of a phosphorus atom can be written


$$1s^22s^22p^63s^23p^3$$

An alternative way to express the electronic configuration is




 $2s^2$









(i) State what is meant by the two arrows in the first box.

(1)

(ii) State why the arrows are all pointing in the same direction in the 3p boxes.

(1)

(Total for Question 1 = 3 marks)

- 2 This question is about ionisation energies.
 - (a) (i) Which equation represents the **second** ionisation of bromine?

(1)


- \square **A** Br(g) + e⁻ \rightarrow Br⁻(g)
- \square **B** Br⁻(g) + e⁻ \rightarrow Br²⁻(g)
- \square **C** Br(g) 2e⁻ \rightarrow Br²⁺(g)
- \square **D** Br⁺(g) e⁻ \rightarrow Br²⁺(g)
- (ii) Which set of successive ionisation energies is most likely to be associated with the element boron?


(1)

- **A** 738, 1451, 7733, 10541, 13629
- **B** 801,2427,3660,25026,32828
- **□ C** 1086,2353,4621, 6223,37832
- **D** 1402,2856,4578, 7475, 9445


(b) (i) Complete the graph to show how the first ionisation energies of the Period 3 elements change across the period. Precise figures are not required.

(3)





(ii) The successive ionisation energies of sodium are shown on the graph.



State what deductions can be made from this graph.

(2)

(Total for Question 2 = 7 marks)

3	Nitrogen forms several hydrides. In addition to ammonia, NH ₃ , it forms hydrazine, N ₂ H ₄
	in which the two nitrogen atoms are covalently bonded together.

(a) (i) Explain what is meant by a covalent bond.

(2)

(ii) Draw a dot-and-cross diagram for hydrazine, showing the outer electrons only.

Use crosses (x) to represent the electrons from nitrogen and dots (•) to represent the electrons from hydrogen.

(1)

(iii) Estimate the H—N—H bond angle in hydrazine.

(1)

Bond angle =

(b) Hydrazine	is very	soluble	in water.
---------------	---------	---------	-----------

Explain, using a labelled diagram and naming the relevant intermolecular interactions, why hydrazine is **very** soluble in water.

(3)

(c) Hydrazine has been used as a rocket fuel.

It is a powerful reducing agent and will react very exothermically with oxidising agents such as hydrogen peroxide.

The equation for the reaction of hydrazine with hydrogen peroxide is

$$N_2H_4(I) \; + \; 2H_2O_2(I) \; \rightarrow \; N_2(g) \; + \; 4H_2O(g)$$

Give **two** reasons why hydrazine is a good rocket fuel when reacted with hydrogen peroxide.

(2)

(Total for Question 3 = 9 marks)



- **4** This question is about isotopes.
 - (a) The table shows data for some isotopes of potassium.

Isotope	Relative isotopic mass	Abundance %
³⁹ K	38.9637	93.218
⁴⁰ K	39.9340	0.012
⁴¹ K	40.9618	6.770

(i)	State what is meant by the terms 'relative isotopic mass' and 'relative atomic ma	ass'.
		(3)
(ii)	State what is meant by the term 'isotopes'. Illustrate your answer by referring to the isotopes of potassium.)
	the isotopes of potassium	(2)


(iii) Use the data in the table to calculate the relative atomic mass of potassium. Give your answer to 4 significant figures.

(2)



(b) The relative isotopic abundances of an element can be measured using a mass spectrometer.

A simplified and incompletely labelled diagram of a mass spectrometer is shown.



(i) Name the feature of the mass spectrometer responsible for the behaviour of the ions in the region indicated by the arrow \mathbf{X} .

(1)

(ii) Explain the three ion pathways, A, B and C, shown in the region indicated by the arrow **X**.

(3)

(1)

(iii) Give a reason why the mass spectrometer must be operated under vacuum.

(Total for Question 4 = 12 marks)



- 5 This question is about the reactions of the halogens and halide ions.
 - (a) (i) When chlorine gas is bubbled through an aqueous solution of potassium iodide, the reaction involves

(1)

- A oxidation only
- Reduction only
- C redox
- D disproportionation
- (ii) Cyclohexane was added to the resulting solution from (a)(i). The mixture was shaken and then allowed to stand for a few minutes. Two layers were formed.

[Density: aqueous layer solution = $1.10 \,\mathrm{g}\,\mathrm{cm}^{-3}$, cyclohexane layer = $0.78 \,\mathrm{g}\,\mathrm{cm}^{-3}$]

The colour of the **lower** layer was

(1)

- A pale yellow
- B purple
- C red
- D pale green
- (b) Concentrated sulfuric acid was added to a small quantity of solid potassium iodide in a test tube.
 - (i) In this exothermic reaction, which of the following mixtures of gases would be produced?

(1)

- A hydrogen iodide and sulfur dioxide only
- B hydrogen iodide and hydrogen sulfide only
- C hydrogen iodide, sulfur dioxide and hydrogen sulfide
- D hydrogen iodide, hydrogen sulfide and iodine

(ii) Hydrogen iodide is a gas which reacts in a similar way to hydrogen chloride.

State the observation when the hydrogen iodide gas is passed over the mouth of an open bottle of concentrated ammonia solution. Write an equation, including state symbols, for the reaction.

(3)

Observation

Equation

(c) Potassium iodate(V) can be prepared by adding solid iodine to a **hot** aqueous solution of potassium hydroxide.

The equation for the reaction is

$$3I_2 + 6KOH \rightarrow KIO_3 + 5KI + 3H_2O$$

Potassium iodate(V) can be separated from the other reaction product using their differing solubilities in water.

	Solubility in water at 25 °C / mol dm ⁻³
KI	8.92
KIO ₃	0.43

(i) Outline a procedure that you could use to obtain a sample of dry, solid potassium iodate(V) from the reaction mixture.

(3)

|
 |
|------|------|------|------|------|------|------|
|
 |
 | |
 | | | |
|
 |
|
 |
|
 |
|
 |
|
 |

(ii) Describe how you would show that iodide ions are present in an aqueous solution of potassium iodide.

(2)

(d) Fluorine is an element in Group 7.

Group 7 includes the elements chlorine, bromine and iodine.

Some information about the melting and boiling temperatures of Group 7 elements is shown in the table.

Element	Melting temperature / K	Boiling temperature / K
chlorine	172	238
bromine	266	332
iodine	387	457

Which is the expected boiling temperature of fluorine, in kelvin, K?

(1)

- **A** 4
- **■ B** 85

(Total for Question 5 = 12 marks)

BLANK PAGE



6	Malachite is a green mineral with the (a) What is the percentage by mass of A 40.3%	formula $Cu_2CO_3(OH)_2$. It has a molar mass of 22 f copper in pure malachite?	1 g mol ⁻¹ .
	■ B 51.4%		
	☑ C 57.5%		
	■ D 67.9%		
	(b) Describe what you would expect t is added to a sample of pure solid	to see when an excess of dilute hydrochloric aci malachite.	d (3)
	(c) (i) Describe how you would carry	out a flame test on a sample of powdered mala	achite. (3)



(ii) When the atoms of some elements are heated, they produce a characteristic flame colour. For example, the copper in malachite gives a blue-green colour. Explain how atoms of different elements can produce different characteristic flame colours when heated.	
	(4)

(d) (i) When malachite is heated to approximately 300 °C, water, carbon dioxide and copper(II) oxide are formed.

The equation for this decomposition is

$$Cu_2CO_3(OH)_2 \rightarrow 2CuO + CO_2 + H_2O$$

Calculate the maximum volume of carbon dioxide that could be produced when 0.810 g of malachite is thermally decomposed.

Assume that the gas is collected at a temperature of 25 $^{\circ}\text{C}$ and 101 kPa pressure.

Give your answer to an appropriate number of significant figures and state the units. [The ideal gas equation is pV = nRT. Gas constant $(R) = 8.31 \,\mathrm{J}\,\mathrm{mol}^{-1}\,\mathrm{K}^{-1}$]

(5)

(ii) The gas was collected in a gas syringe with a stated accuracy of $\pm 0.5\,\text{cm}^3$.

Calculate the percentage uncertainty in the volume of gas collected.

(1)



(iii) Malachite ore is a mixture of malachite and rock. A 0.810 g sample of malachite ore was thermally decomposed, producing 0.571 g of copper(II) oxide.

Calculate the percentage purity of this malachite ore sample. Give your answer to an appropriate number of significant figures.

(3)

(Total for Question 6 = 20 marks)



	Specific values of atomic ra			atomic	radii of the	e Perioc	d 2 and	Period 3	3 elemen
	Specific values of atomic for	adii aic	Hotree	funcu.					(3)
								•••••	
,									
⁽ (b)	The melting temperatures	of the	Period 2	2 eleme	nts are sho	own.			
	Symbol of the element	Li	Be	В	C _(diamond)	N	0	F	Ne
	Melting temperature / K	454	1551	2573	3970	63	55	53	25
	Merting temperature / K	757	1331	2373	3370	05] 33	23
	Explain the trend in meltin	ng temp	perature	s acros	s the elem	ents of	Period ¹	2 in	
						CIICS OI	i ciioa i	<u> </u>	
	terms of their structure an	d bond				C1115 01	r criou .	2 111	(6)
	terms of their structure an	d bond					r crioa .	2 111	(6)
	terms of their structure an	d bond							(6)
	terms of their structure an	d bond					T CHOO!		(6)
	terms of their structure an	d bond							(6)
	terms of their structure an	d bond							(6)
	terms of their structure an	d bond							(6)
	terms of their structure an	d bond							(6)
	terms of their structure an	d bond							(6)
	terms of their structure an	d bond							(6)
	terms of their structure an	d bond							(6)
	terms of their structure an	d bond							(6)
	terms of their structure an	d bond							(6)
	terms of their structure an	d bond							(6)
	terms of their structure an	d bond							(6)

 (Total for Question 7 = 9 marks)

- 8 This question is about the thermal stability of Group 1 and Group 2 nitrates and carbonates.
 - (a) Complete the equations for the thermal decomposition of sodium nitrate, $NaNO_3$, and for the thermal decomposition of calcium nitrate, $Ca(NO_3)_2$. State symbols are not required.

(2)

$$NaNO_3 \rightarrow$$

$$Ca(NO_3)_2 \rightarrow$$

(b) The thermal stability of Group 1 nitrates increases down the group.

The decomposition temperatures of some Group 1 nitrates are shown.

Name	Formula	Decomposition temperature / K
sodium nitrate	NaNO ₃	653
potassium nitrate	KNO ₃	673
caesium nitrate	CsNO ₃	687

Explain why the thermal stability of caesium nitrate is greater than that of sodium nitrate.

(3)

20

TOTAL FOR PAPER = 80 MARKS						
(Total for Question 8 = 8 ma	rks)					
(iii) Give a reason, in terms of the bonding, why a high decomposition temperature is required.	(1)					
(ii) Name all the types of bond present in calcium carbonate.	(1)					
(c) Calcium carbonate is thermally decomposed during the manufacture of cement.(i) Write an equation, including state symbols, for the thermal decomposition of calcium carbonate.	(1)					



BLANK PAGE



BLANK PAGE



	7 0 (8)	Helium (17)	19.0 F	9 35 5	CI Ar chlorine argon 17 18	6.62	_	bromine krypton 35 36	126.9 131.3		53 54	[210]	Ą	astatine 85	The second secon	Elements with atomic numbers 112-116 have been reported but not fully authenticated		175	Ξ	lutetium 71	[257]	Lr lawrencium
	0	(16)	16.0 O oxygen	8 22	Sulfor 16	0.67	Se	selenium 34	127.6	Тe	tellurium 52	[506]	8	polonium 84		nticated		173	ХÞ	ytterbium 70	[254]	Nobelium
u	n	(15)	14.0 N	21.0	P phosphorus 15	74.9	As	arsenic 33	121.8	Sb	antimony 51	209.0	<u>B</u>	bismuth 83		but not fully authenticated		169	٤	thulium 69	[326]	Md
,	4	(14)	12.0 C carbon	78.1		72.6	g	germanium 32	118.7	Sn	20 E	207.2	Ъ	lead 82		but not f		167	ដ	erbium 68	[253]	Fm fermium
•	7	(13)	10.8 B boron	27.0	AI aluminium 13	2'69	g	gallium 31	114.8	占	indium 49	204.4	F	thallium 81	140	nents with		165		holmium 67	[254]	Cf Es californium einsteinium
					(12)	65.4	Zu	zinc 30	112.4	8	cadmium 48	200.6	Hg	mercury 80				163	ò	dysprosium 66	[251]	Cf californium
			(11)				D U	copper 29	107.9	Ag	silver 47	197.0	Αn	gold 79	[272]	roentgenium	111	159	4	terbium 65	[245]	BK berkelium
en la	(01)				58.7	ž	nicket 28	106.4	Pd	palladium 46	195.1	£	platinum 78	[271]	darmstadfium	110	157	В	gadolinium 64	[247]	Carricum	
1.0 H hydrogen 1			(6)		58.9	ဝ	cobalt 27	102.9	R	rhodium 45	192.2	1	iridium 77	[268]	Mit	109	152	En	europium 63	[243]	Am americium	
		1.0 H hydrogen		(8)	55.8	Fe	iron 26	1.101	Ru	ruthenium 44	190.2	os	osmium 76	[777]	hassium	108	150		samarium 62	[242]	Pu plutonium	
		6						manganese 25	[86]		technetium 43	186.2	Re	rhenium 75	_	bohrium		[147]	Рш	promethium 61	[237]	Np neptunium
			mass	umber	(9)	52.0	ъ	E	626	Wo	molybdenum 42	183.8	>	tungsten 74	[596]	Seaborgium	106	144	P	neodymium 60	238	U
		Key	relative atomic mass atomic symbol	atomic (proton) number	(5)	50.9	>	vanadium 23	67.6	Ą	niobium 41	180.9	ā	tantalum 73	_	dubníum		141	Pr	praseodymium neodymium promethium 59 60 61	[231]	Pa protactinium
			relat	atomic	9	47.9	j드	titanium 22	91.2	JΖ	zirconium 40	178.5	ŧ	hafnium 72	[261]	rutherfordium	104	140	e ئ	cerium 58	232	thorium
	(3)					45.0	Sc aandium 21 88.9 Y Y Y 4ttrium 39 138.9 La* nthanum 57 [227] Ac* ctinium 89				Se											
•	7	(2)	9.0 Be beryllium	74.2	Mg magnesium 12	40.1	S	calcium 20	97.8	'n	strontium 38	137.3		barium 56	[977]	radium	88		* Lanthanide series	* Actinide series		
	-	ε	6.9 Li lithium	3	Na Sodium	39.1	¥	potassium 19	85.5	8	75 37	132.9	S	caesium 55	[223]	francium	87		. Lanth	* Actini		